

## Chromosome reconfigurations

These were once known as “chromosomal aberrations.” They are now recognized as intermediate steps that take place as the genome reconfigures itself.

### Deficiencies

First described by Bridges in the notched wing mutants of *Drosophila*

Later, Mohr was first to correlate gametic loss with a cytological loss




Figure 1, <https://www.slideshare.net/Samchuchoo/chromosomal-aberration>

#### McClintock, 1931

Correlated genetic loss with pachytene analysis:

- Cross female *pl/pl* × X-rayed male *Pl/Pl* → looked for F1 plants of *pl* phenotype, then did pachytene analysis
- Pseudodominance: the expression of a recessive phenotype due to a deficiency
- Other phenotypic effects
  - Male sterility (lethal in male gamete)
  - Reduced female transmission




Figure 2. A: Interstitial deficiency of chromosome 7. B: Terminal deficiency of chromosome 6 (McClintock, 1931)

#### McClintock, 1944


Working with the *yg2* (yellow-green) locus, which was known to be 15 crossover units from *C* (colored endosperm) on chromosome 9

She also found *py* (pale-yellow) mutants and *w* (white) mutants that mapped to the same region. In the following crosses:

- *py* × *w* → all pale-yellow, therefore, were allelic, with *py* dominant over *w*
- *w* × *yg2* → all yellow-green, therefore, were allelic
- *py* × *yg2* → green, therefore, were not allelic, which is a discrepancy

This genetic discrepancy can be explained by cytological deficiencies:

- Normal chromosome 9, with *Yg2*
- Normal chromosome 9, with *yg2*
- Chromosome 9 missing terminal knob, but with the wild-type allele (*Yg2*)
- Chromosome 9 missing the *Yg2* locus altogether



Thus:  $b + b =$  yellow-green seedling       $b + c =$  green seedling  
 $c + c =$  pale-yellow seedling       $b + d =$  yellow-green seedling  
 $d + d =$  white seedling       $c + d =$  pale-yellow seedling

- Deficiencies in plants not as frequently transmitted as in animals, as haploid gametophyte will not tolerate significant deficiencies
- Recover more aberrations through the female than the male, as the ovule around the female gametophyte serves as nurse tissue
- Alternative hypothesis: Less genes are turned on in the female
- Diploids cannot tolerate homozygous deficiencies above a certain size. Allopolyploids can.

### 5A deficiencies in wheat and oat ( $2n = 6x = 42$ ):

Occasionally get off-types that resemble related wild species

- Due to addition/deletion of chromosome 5A


Oat	Wheat	
Fatoid	Speltoid	
	Het speltoid	
	Subnormal	
Steriloid	Subcompactoid	
	Compactoid	



*Speltoid, normal, subcompactoid, and compactoid heads of wheat*

## Mapping deficiencies

In the example below, the chromosome carrying the "A" allele is not transmissible through a gamete because of the deficiency:




However, "A" can still be recovered if there is a crossover between "A" and the deficiency, allowing "A" to be mapped relative to the junction. E.g.:

- If out of 179 progenies, 146 are "a" and 33 are "A", then the distance between "A" and the junction is:

$$\frac{A}{a+A} = \frac{33}{179} = 0.184$$

- which = the % of dominant progeny
- In this case, every CO gives 1 dominant progeny, as compared to the normal situation where 1 CO gives a 50% recombination rate.

## Effect of deficiencies on linkage maps



If the deficiency is viable, a homozygous map would simply be missing all the loci in the deficiency, while a heterozygous map would have all the loci in the deficient region map to one point.

## Uses of deficiencies

- Make knockouts
- Cytological mapping via pseudodominance- **Khush & Rick, 1968**
- Studies of pairing specificity
- Gene cloning through genomic subtraction
- Reverse genetics
- Determine function of the chromosomal region

- Fine genetic mapping of mutations and markers
  - Eg., Merlino et al., 2012 - Associate loss of proteins from 2D gels with specific deletions
  - Missing protein meant the deletion has the coding gene or a regulatory gene




Merlino et al., 2012

## Detection

### Comparative Genomic Hybridization (CGH)

Makes it much easier to ID region where deficiency is, and thus assign phenotype to gene.

An array is made of single copy DNA in linear order as is found in the chromosomes



- Reference and deletion genomes are both labeled
- When DNA from both is present, the labels cancel each other out
- If a deletion is present, there is no counter label, making the deletion obvious

## Structural variants: Indels

It is also possible to get smaller indels.

- These do not affect fertility, but can affect the phenotype
- 3 examples are:

### Non-pungent chili peppers


Stewart et al., 2005; Hulse-Kemp et al., 2018



- 2570 pb deletion
- Eliminates hotness


### Tomato fruit size

Mu et al., 2017





### Determinancy in bean

Kwak et al., 2012



### Ionizing radiation creates deficiencies

Although Muller is recognized as the discoverer of radiation-induced mutations in 1927, for which he received the Nobel Prize, radiation mutagenesis traces back to Emmy Stein in 1921



## Structural variants: Mutation breeding

When the needed mutation is not found in nature, it can be obtained by inducing structural variation, usually with ionizing radiations, which induce:

- Indels 4pb - 8kb
- Inversions up to 1.5 kb
- Frame shift mutations
- Premature stop codons






Figure 3. Institute of Radiation Breeding, Ibaraki-ken, JAPAN

## Duplications

Types: Tandem, Reversed, & Displaced

Displaced can be:

- Fragment with centromere
- To another chromosome
- To another location on the same chromosome



Nature Education Adapted from Pierce, Benjamin. *Genetics: A Conceptual Approach, 2nd ed*

Modified from Randall, PJ. *iGenetics*, Pearson Education Inc, published as Benjamin Cummings


- First found by Bridges in *Drosophila*

## Behavior of duplications


### Bridges 1936

Bar eye, a sex-linked dominant trait

- In female flies, bar would mutate to double bar or back to normal
- Studying salivary gland chromosomes, found that bar was a duplication of bands 1-7 in the 16A chromosomal region.



## Duplications prone to unequal crossover



Modified from <https://slideplayer.com/slide/10755099/>

- This illustrates one of the principle properties of duplications -- their propensity for unequal crossing over
- Duplications provide a way of adding genetic material
- Increase copy number of a gene
- 2 genes, identical by duplication, can then diverge by mutation to achieve different functions

## Uses

Study effect of gene dosage


Increase amount of desired gene product without resorting to polyploidy

## Structural variants: Duplications

Structural variant duplications – affect the phenotype, but too small to affect fertility. They are prone to rearrangements that affect the phenotype (Sub 1 rice; Sorghum) or trigger RNAi (soybean, maize, rice examples)

### Sub1 rice


Xu et al., 2006



Soybean Vodkin seed coat color eg is missing here and in handout


### Sorghum dwarfism

Multani et al., 2003



### C-1df green stalks in corn

Della Vedova et al., 2005



### Low glutelin rice

Kusaba et al., 2003

